

# وبینار علمی مراقبت های پرستاری در تروما

chest wall trauma

تهیه و تنظیم:

پرستو پورجعفری مقدم

کارشناسی ارشد پرستاری مراقبت های ویژه

بیمارستان آریا


مهر 1400

# Introduction




# **Introduction:**

- ❑ Traumatic injuries to the chest contribute to 75% of all traumatic deaths.
- ❑ Thoracic injuries range from simple rib fractures to complex life-threatening rupture of organs.
- ❑ The mechanisms of injuries causing chest trauma are separated into two categories: blunt trauma and penetrating trauma.

- 
- ❑ Chest injuries are potentially life-threatening because of immediate disturbances of cardiorespiratory physiology and haemorrhage and later developments of infection, damaged lung and thoracic cage.





## **Incidence:**



- 25% of all death form traumatic injury.



```
graph TD; A["Causes:"] --> B["BLUNT INJURY CAUSES"]; A --> C["PENETRATING INJURY CAUSES"]
```

## Causes:

**BLUNT INJURY  
CAUSES**

**PENETRATING  
INJURY CAUSES**







## Blunt Trauma to the Chest









## ❖ BLUNT INJURY CAUSES:

- ❑ Motor vehicle accident
- ❑ Pedestrian accident
- ❑ Fall
- ❑ Sports injury
- ❑ Assault with blunt object or Altercations
- ❑ Crush injury
- ❑ Explosion



## ❖ PENETRATING INJURY CAUSES:

- ❑ Knife
- ❑ Gunshot
- ❑ Stick
- ❑ Arrow
- ❑ Occupational injury




## Rib Fracture:

---

- Most common chest injury.
- May interfere with ventilation and may lacerate underlying lung.







# Hemothorax:


---

- ❑ Blood in pleural space as a result of penetrating or blunt chest trauma.
- ❑ Accompanies a high percentage of chest injuries.
- ❑ Can result in hidden blood loss.


Blood-filled  
pleural space

Lung

B




# Hemothorax



## Flail Chest:

---

- ❑ Loss of stability of chest wall as a result of multiple rib fractures, or combined rib and sternum fractures.
- ❑ When this occurs, one portion of the chest has lost its bony connection to the rest of the rib cage.




# Pulmonary Contusion:

---

- Bruise of the lung parenchyma those results in leakage of blood and edema fluid into the alveolar and interstitial spaces of the lung.
- May not be fully developed for 24 to 72 hours.

## ██████████: Thoracic Injuries




# Cardiac Tamponade:

---


- Compression of the heart as a result of accumulation of fluid within the pericardial space.
- Caused by penetrating injuries, metastasis, and other disorders.



Healthy



Tamponade



# Clinical manifestation:



# Respiratory


---

- ❑ Dyspnea, respiratory distress
- ❑ Cough with or without haemoptysis
- ❑ Cyanosis of mouth, face, nail beds, mucous membranes
- ❑ Tracheal deviation
- ❑ Audible air escaping from chest wound
- ❑ Decreased breath sounds on side of injury
- ❑ Decreased O<sub>2</sub> saturation
- ❑ Frothy secretions

# Cardiovascular

---

- ❑ Rapid, thready pulse
- ❑ Decreased BP
- ❑ Narrowed pulse pressure
- ❑ Asymmetric BP values in arms
- ❑ Distended neck veins
- ❑ Muffled heart sounds

- 
- ❑ Chest pain
  - ❑ Crunching sound synchronous with heart sounds
  - ❑ Dysrhythmias

# Surface Findings

---

- ❑ Bruising
- ❑ Abrasions
- ❑ Open chest wound
- ❑ Asymmetric chest movement
- ❑ Subcutaneous emphysema

100





- **History collection**

- **Physical examination**


- While doing physical examination assess for abdominal tenderness, chest tenderness, chest bruising, chest swelling, decrease lung sound, wheezing, rapid pulse and rapid breathing, chest crepitation, cyanosis, dyspnea.

- X- Ray

- CT Scan and MRI





- 
- The **goal** is to restore normal **cardiorespiratory** function as quickly as possible.
  - This is accomplished by,
    - Performing effective resuscitation
    - While simultaneously assessing the patient,
    - Restoring chest wall integrity,
    - Reexpanding the lung.

## Rib Fracture:

- Give analgesics (usually nonopioid) to assist in effective coughing and deep breathing.
- Encourage deep breathing with strong inspiration; give local support to injured area by splinting with hands.

# Hemothorax:

- Assist with **thoracentesis** to aspirate blood from pleural space, if being done before a chest tube insertion.
- Assist with **chest tube insertion** and set up drainage system for complete and continuous removal of blood and air.
  - ▣ Auscultate lungs and monitor for relief of dyspnea.
  - ▣ Monitor amount of blood loss in drainage.
- Replace volume with I.V. fluids or blood products.



## Flail Chest:

---

- ❑ Stabilize the flail portion of the chest with hands; apply a pressure dressing and turn the patient on injured side, or place 10-lb sandbag at site of flail.
- ❑ Thoracic epidural analgesia may be used for some patients to relieve pain and improve ventilation.



# Pulmonary Contusion:

---



- Employ mechanical ventilation to keep lungs inflated.
- Administer diuretics to reduce edema.
- Correct metabolic acidosis with I.V. sodium bicarbonate.
- Use PAP monitoring.
- Monitor for development of pneumonia.

# Cardiac Tamponade:

---

- Assist with **pericardiocentesis** to provide emergency relief and improve hemodynamic function until surgery can be undertaken.
- Prepare for emergency **thoracotomy** to control bleeding and to repair cardiac injury.








Source: J.E. Tintinalli, J.S. Stapczynski, O.J. Ma, D.M. Yealy, G.D. Meckler, D.M. Cline:  
 Tintinalli's Emergency Medicine: A Comprehensive Study Guide, 8th Edition  
[www.accessmedicine.com](http://www.accessmedicine.com)  
 Copyright © McGraw-Hill Education. All rights reserved.


## **ADDITIONAL RESPONSIBILITIES:**

- Secure and support the airway as indicated.

- 
- Prepare for tracheostomy if indicated.
    - ▣ Tracheostomy helps to clear tracheobronchial tree, helps the patient breathe with less effort, decreases the amount of dead airspace in the respiratory tree, and helps reduce paradoxical motion.
    - ▣ When used with mechanical ventilation, provides a closed system and stabilizes the chest.

- 
- ❑ Secure one or more I.V. lines for fluid replacement, and obtain blood for baseline studies, such as hemoglobin level and hematocrit.
  - ❑ Monitor serial CVP readings to prevent hypovolemia and circulatory overload.
  - ❑ Monitor ABG/Spo<sub>2</sub> results to determine need for supplemental oxygen, mechanical ventilation.

- 
- Institute ECG monitoring for early detection and treatment of cardiac dysrhythmias (dysrhythmias are a frequent cause of death in chest trauma).

- 
- Maintain ongoing surveillance for complications:
    - ▣ Aspiration
    - ▣ Atelectasis
    - ▣ Pneumonia
    - ▣ Mediastinal/subcutaneous emphysema
    - ▣ Respiratory failure

# **Patient education and health maintenance:**

---

- Instruct patient in splinting techniques.
- Make sure patient is aware of importance of automobile seat belt use to reduce serious chest injuries caused by automobile accidents.
- Teach patient to report signs of complications increasing dyspnea, fever, and cough.



